

Chapter 3

Exponential and Logarithmic Functions

3.3 Properties of Logarithms

Chapter 3

Homework

Chapter 3.3

Objectives

- ☐ Use the product rule.
- ☐ Use the quotient rule.
- ☐ Use the power rule.
- ☐ Expand logarithmic expressions.
- ☐ Condense logarithmic expressions.
- ☐ Use the change-of-base property.

Why Logarithms

■ Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

- Large numbers are difficult for people to assimilate. Donald Trump claims to be worth in excess of \$10 billion, the U.S. budget debt is in the trillions of dollars, but what do those numbers represent?
- Well a million seconds is about 11.5 days, a billion seconds is about 31 years, and one trillion seconds is about 31,710 years. That is the difference between a year's vacation time, the time it takes to get the kids out of the house, and the entirety of human existence.
- To change these numbers into manageable size we use logarithms. The log of 1,000,000 is 6, $\log 1,000,000,000 = 9$, and $\log 1,000,000,000,000 = 12$. 6, 9, 12 are much easier to handle. We accomplish this through the use of **exponents**.

The Product Rule

Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

Let b , M , and N be positive real numbers with $b \neq 1$. Then:

$$\log_b(MN) = \log_b M + \log_b N$$

The logarithm of a **product** is the **sum** of the logarithms.

This can be easily verified by converting to exponential form.

Let $\log_b(M) = x$ $\log_b(N) = y$ Then $b^x = M$ $b^y = N$

$$\log_b(MN) = \log_b(b^x \cdot b^y) = \log_b(b^{x+y}) = x + y = \log_b M + \log_b N$$

Example: Using the Product Rule

Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

Use the product rule to expand each logarithmic expression:

$$\log_6(7 \cdot 11) = \log_6 7 + \log_6 11$$

$$\log 100x = \log 100 + \log x = 2 + \log x$$

$$\log_6 648 = \log_6(216 \cdot 3) = \log_6 216 + \log_6 3 = 3 + \log_6 3$$

Solve for y : $\ln y = \ln x + \ln c = \ln(x \cdot c)$

$$y = x \cdot c$$

$$\log_b(MN) = \log_b M + \log_b N$$

The Quotient Rule

- Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

- Let b , M , and N be positive real numbers with $b \neq 1$. Then:

$$\log_b \left(\frac{M}{N} \right) = \log_b M - \log_b N$$

- The logarithm of a **quotient** is the **difference** of the logarithms.
- This is simply a rewrite of the previous rule, remembering subtraction is addition of the opposite.

Example: Using the Quotient Rule

Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

Use the quotient rule to expand each logarithmic expression:

$$\log_8 \left(\frac{23}{x} \right) = \log_8 23 - \log_8 x$$

$$\begin{aligned} \ln \left(\frac{e^5}{11} \right) &= \ln e^5 - \ln 11 \\ &= 5 - \ln 11 \end{aligned}$$

$$\log \left(\frac{\sqrt{x} \sqrt[3]{y^2}}{z^4} \right) = \log \sqrt{x} + \log \sqrt[3]{y^2} - \log z^4$$

$$\log_b \left(\frac{M}{N} \right) = \log_b M - \log_b N$$

We will finish this in a couple of slides.

The Power Rule

Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

Let b , M , and N be positive real numbers with $b \neq 1$. Then:

$$\log_b M^p = p \log_b M$$

The logarithm of a number with an exponent is the product of the exponent and the logarithm of that number.

This can again be shown by converting to exponential form.

Let $\log_b M = a$ Then $b^a = M$

$$M^p = (b^a)^p = b^{pa} \quad \log_b M^p = pa = p \log_b M$$

Using the Power Rule

Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

Use the power rule to expand each logarithmic expression:

$$\log_b M^p = p \log_b M$$

$$\log_6 3^9 = 9 \log_6 3$$

$$\ln \sqrt[3]{x} = \ln x^{\frac{1}{3}} = \frac{1}{3} \ln x$$

$$\log(x+4)^2 = 2 \log(x+4)$$

$$\log_{25} 5^3 = 3 \log_{25} 5 = 3 \log_{25} 25^{\frac{1}{2}} = \frac{3}{2} \log_{25} 25 = \frac{3}{2}$$

$$\begin{aligned} \log \left(\frac{\sqrt{x} \sqrt[3]{y^2}}{z^4} \right) &= \log \sqrt{x} + \log \sqrt[3]{y^2} - \log z^4 \\ &= \log x^{\frac{1}{2}} + \log (y^2)^{\frac{1}{3}} - \log z^4 \\ &= \frac{1}{2} \log x + \frac{2}{3} \log y - 4 \log z \end{aligned}$$

Properties for Expanding Logarithmic Expressions

- Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

For $M > 0$ and $N > 0$:

Product Rule

$$\log_b(MN) = \log_b M + \log_b N$$

Quotient Rule

$$\log_b \left(\frac{M}{N} \right) = \log_b M - \log_b N$$

Power Rule

$$\log_b M^p = p \log_b M$$

Note: In every case the base remains consistent (the same).

Common Errors to Avoid

- Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

$$\log_b(M+N) \neq \log_b M + \log_b N$$

$$\log_b(MN) \neq \log_b M \bullet \log_b N$$

$$\log_b\left(\frac{M}{N}\right) \neq \frac{\log_b M}{\log_b N}$$

$$\log_b M^p \neq \log_b pM$$

$$\log_b MN^p \neq p \log_b MN$$



Expanding Logarithmic Expressions

Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

Expand the logarithmic expression as much as possible:

$$\begin{aligned}\log_b \left(x^4 \sqrt[3]{y} \right) &= \log_b \left(x^4 y^{\frac{1}{3}} \right) \\ &= \log_b \left(x^4 \right) + \log_b \left(y^{\frac{1}{3}} \right) \\ &= 4 \log_b x + \frac{1}{3} \log_b y\end{aligned}$$

Ejemplo

■ Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

● Write the following expression in terms of logs of x and z .

$$\begin{aligned}\log\left(x\sqrt{\frac{\sqrt{x}}{z}}\right) &= \log x + \log\sqrt{\frac{\sqrt{x}}{z}} \\ &= \log x + \log\left(\frac{\sqrt{x}}{z}\right)^{\frac{1}{2}} \\ &= \log x + \frac{1}{2}\log\left(\frac{\sqrt{x}}{z}\right)\end{aligned}$$

$$\begin{aligned}&= \log x + \frac{1}{2}\left(\log\sqrt{x} - \log z\right) \\ &= \log x + \frac{1}{2}\left(\log x^{\frac{1}{2}} - \log z\right) \\ &= \log x + \frac{1}{2}\left(\frac{1}{2}\log x - \log z\right) \\ &= \log x + \frac{1}{4}\log x - \frac{1}{2}\log z\end{aligned}$$

Expanding Logarithmic Expressions

■ Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

● Use the product rule to expand each logarithmic expression as much as possible:

$$\begin{aligned}\log_5 \left(\frac{\sqrt{x}}{25y^3} \right) &= \log_5 \left(\frac{x^{\frac{1}{2}}}{25y^3} \right) = \log_5 x^{\frac{1}{2}} - \log_5 25y^3 \\ &= \log_5 x^{\frac{1}{2}} - (\log_5 25 + \log_5 y^3) \\ &= \frac{1}{2} \log_5 x - (\log_5 25 + 3 \log_5 y) \\ &= \frac{1}{2} \log_5 x - 2 - 3 \log_5 y\end{aligned}$$

Expanding Logarithmic Equation

■ Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

● Write the following expression in terms of logs of x , y , and z .

$$\begin{aligned}\log\left(\sqrt{\frac{xy^2}{z^8}}\right) &= \log\left(\frac{xy^2}{z^8}\right)^{\frac{1}{2}} &&= \frac{1}{2}(\log x + \log y^2 - \log z^8) \\ &= \frac{1}{2}\log\left(\frac{xy^2}{z^8}\right) &&= \frac{1}{2}(\log x + 2\log y - 8\log z) \\ &= \frac{1}{2}(\log(xy^2) - \log z^8) &&= \frac{1}{2}\log x + \log y - 4\log z\end{aligned}$$

Condensing Logarithmic Expressions

- Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

For $M > 0$ and $N > 0$:

Product Rule

$$\log_b M + \log_b N = \log_b (MN)$$

Quotient Rule

$$\log_b M - \log_b N = \log_b \left(\frac{M}{N} \right)$$

Power Rule

$$p \log_b M = \log_b M^p$$

Note: In every case the base remains consistent (the same).

Condensing Logarithmic Expressions

Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

Write as a single logarithm

$$\log 25 + \log 4 = \log(25 \cdot 4) = \log 100 = 2$$

$$\log(7x + 6) - \log x = \log\left(\frac{7x + 6}{x}\right)$$

$$\begin{aligned} 2\ln x + \frac{1}{3}\ln(x + 5) &= \ln x^2 + \ln(x + 5)^{\frac{1}{3}} \\ &= \ln x^2 + \ln \sqrt[3]{x + 5} \\ &= \ln x^2 \sqrt[3]{x + 5} \end{aligned}$$

Solving Logarithmic Equation

■ Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

● Find x if $2\log_b 5 + \frac{1}{2}\log_b 9 - \log_b 3 = \log_b x$

$$\log_b 5^2 + \log_b 9^{\frac{1}{2}} - \log_b 3 = \log_b x$$

$$\log_b 25 + \log_b 3 - \log_b 3 = \log_b x \quad \longrightarrow \quad x = 25$$

● or, if you prefer $\log_b \left(\frac{25 \cdot 3}{3} \right) = \log_b x$

The Change-of-Base Property

- Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

- For any logarithmic bases a and b , and any positive number M ,

$$\log_b M = \frac{\log_a M}{\log_a b}$$

- The logarithm of M with base b is equal to the logarithm of M with any new base divided by the logarithm of b with that new base.
- The change of base formula allows you to estimate any log on your calculator without using `logbase()`, or to use a simpler calculator (i.e. TI 30XIIS).

The Change-of-Base Property

- Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

Change-of-Base Formula

Let a , b , and x be positive real numbers such that $a \neq 1$ and $b \neq 1$. Then $\log_a x$ can be converted to a different base as follows.

Base b

$$\log_a x = \frac{\log_b x}{\log_b a}$$

Base 10

$$\log_a x = \frac{\log x}{\log a}$$

Base e

$$\log_a x = \frac{\ln x}{\ln a}$$

The Change-of-Base and Common and Natural Logarithms

- Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

- Introducing Common Logarithms

$$\log_b M = \frac{\log M}{\log b}$$

- Introducing Natural Logarithms

$$\log_b M = \frac{\ln M}{\ln b}$$

Changing Base to Common Logarithms

- Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

- Use your calculator and common logs to evaluate

$$\log_7 2506 = \frac{\log 2506}{\log 7} \approx \frac{3.39898}{.8451} \approx 4.022$$

- Use your calculator and natural logs to evaluate

$$\log_7 2506 = \frac{\ln 2506}{\ln 7} \approx \frac{7.8264}{1.9489} \approx 4.022$$

Finding the inverse of a function

■ Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

● Remember, when finding the inverse of a function:

1. Replace $f(x)$ with y in the equation.
2. Exchange x and y in the equation.
3. Solve the equation for y in terms of x .
4. If the inverse is a function, replace y with $f^{-1}(x)$.

● These same rules apply when finding the inverse of an exponential equation (a log) or when finding the inverse of a log function (an exponential function).

Finding the Inverse of an Exponential Equation.

Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

Find the inverse of $f(x) = 2 \cdot 5^x - 3$

1. Replace $f(x)$ with y in the equation. $y = 2 \cdot 5^x - 3$

2. Exchange x and y in the equation. $x = 2 \cdot 5^y - 3$

3. Solve the equation for y in terms of x . $x + 3 = 2 \cdot 5^y$ $\frac{x+3}{2} = 5^y$ $\log_5 \left(\frac{x+3}{2} \right) = y$

4. If the inverse is a function, replace y with $f^{-1}(x)$. $f^{-1}(x) = \log_5 \left(\frac{x+3}{2} \right)$

Finding the Inverse of an Exponential Equation.

- Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

- We can graph both functions


$$f(x) = 2 \cdot 5^x - 3$$

$$f^{-1}(x) = \log_5 \left(\frac{x+3}{2} \right)$$

x	f(x)
-2	-2.92
-1	-2.6
0	-1
1	7
2	47

x	f ⁻¹ (x)
-3	oops
-2	-0.4307
-1	0
0	0.2519
2	0.5693
7	1

- Note asymptotes, intercepts, and axis of symmetry



Finding Asymptotes and Intercepts

Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

$$f(x) = 2 \cdot 5^x - 3$$


$$0 = 2 \cdot 5^x - 3 \quad 3 = 2 \cdot 5^x \quad \frac{3}{2} = 5^x \quad \log_5 \frac{3}{2} = x \approx .25$$

$$y = 2 \cdot 5^0 - 3 \quad y = 2 - 3 = -1$$

$$f^{-1}(x) = \log_5 \left(\frac{x+3}{2} \right)$$

$$0 = \log_5 \left(\frac{x+3}{2} \right) \quad \frac{x+3}{2} = 1 \quad x = -1$$

$$y = \log_5 \left(\frac{0+3}{2} \right) \quad y = \log_5 \left(\frac{3}{2} \right) \approx .2519$$



Finding the Inverse of an Exponential Equation.

Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

Find the inverse of $f(x) = \frac{1}{3}e^x + 1$

1. Replace $f(x)$ with y in the equation. $y = \frac{1}{3}e^x + 1$

2. Exchange x and y in the equation. $x = \frac{1}{3}e^y + 1$

3. Solve the equation for y in terms of x . $x - 1 = \frac{1}{3}e^y$ $3(x - 1) = e^y$ $\ln e^y = \ln(3x - 3)$
 $y = \ln(3x - 3)$

4. If the inverse is a function, replace y with $f^{-1}(x)$. $f^{-1}(x) = \ln(3x - 3)$

Finding the Inverse of an Exponential Equation.

- Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

- We can graph both functions


$$f(x) = \frac{1}{3}e^x + 1$$

$$f^{-1}(x) = \ln(3x - 3)$$

x	f(x)
-2	1.0451
-1	1.1226
0	1.333
1	1.9061
2	3.463

x	f ⁻¹ (x)
1	oops
2	1.0986
3	1.79
4	2.1972
6	2.7081

- Note asymptotes, intercepts, and axis of symmetry



Finding the Inverse of an Exponential Equation.

- Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

$$f(x) = \frac{1}{3}e^x + 1$$


$$0 = \frac{1}{3}e^x + 1 \quad \frac{1}{3}e^x = -1$$

$$y = \frac{1}{3}e^0 + 1 \quad y = \frac{4}{3}$$

$$f^{-1}(x) = \ln(3x - 3)$$

$$0 = \ln(3x - 3) \quad 3x - 3 = 1 \quad x = \frac{4}{3}$$

$$y = \ln(0 - 3)$$



Finding the Inverse of a Logarithm Function

Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

Find the inverse of $f(x) = 2\log_3(x-2) - 1$

1. Replace $f(x)$ with y in the equation. $y = 2\log_3(x-2) - 1$

2. Exchange x and y in the equation. $x = 2\log_3(y-2) - 1$

3. Solve the equation for y in terms of x . $\frac{x+1}{2} = \log_3(y-2)$ $3^{\left(\frac{x+1}{2}\right)} = y-2$ $y = 3^{\left(\frac{1}{2}(x+1)\right)} + 2$

4. If the inverse is a function, replace y with $f^{-1}(x)$. $f^{-1}(x) = 3^{\left(\frac{1}{2}(x+1)\right)} + 2$

Finding the Inverse of a Logarithm Function

Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

● We have graphed enough, so let us find the intercepts.

$$f(x) = 2\log_3(x-2) - 1$$

$$0 = 2\log_3(x-2) - 1 \quad \frac{1}{2} = \log_3(x-2) \quad x = 3^{\frac{1}{2}} + 2 = 2 + \sqrt{3} \approx 3.732$$

$$y = 2\log_3(0-2) - 1$$

$$f^{-1}(x) = 3^{\left(\frac{1}{2}(x+1)\right)} + 2$$

$$0 = 3^{\left(\frac{1}{2}(x+1)\right)} + 2 \quad -2 = 3^{\left(\frac{1}{2}(x+1)\right)}$$

$$y = 3^{\left(\frac{1}{2}(0+1)\right)} + 2 \quad y = 3^{\left(\frac{1}{2}(0+1)\right)} + 2 = 3^{\left(\frac{1}{2}\right)} + 2$$

Finding the Inverse of a Logarithmic Function

Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

Find the inverse of $f(x) = 2\log(2x + 3) - 5$

1. Replace $f(x)$ with y in the equation. $y = 2\log(2x + 3) - 5$

2. Exchange x and y in the equation. $x = 2\log(2y + 3) - 5$

3. Solve the equation for y in terms of x . $\frac{x+5}{2} = \log(2y + 3)$ $10^{\left(\frac{x+5}{2}\right)} = 2y + 3$ $y = \frac{10^{\left(\frac{x+5}{2}\right)} - 3}{2}$

4. If the inverse is a function, replace y with $f^{-1}(x)$. $f^{-1}(x) = \frac{10^{\left(\frac{x+5}{2}\right)} - 3}{2}$

Finding Inverse of Logarithm

■ Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

● Again, find the intercepts.

$$f(x) = 2\log(2x + 3) - 5$$

$$0 = 2\log(2x + 3) - 5 \quad \frac{5}{2} = \log(2x + 3) \quad 10^{\frac{5}{2}} = 2x + 3 \quad x = \frac{10^{\frac{5}{2}} - 3}{2} \approx 156.61$$

$$y = 2\log(2 \cdot 0 + 3) - 5 \quad y = 2\log(3) - 5 \approx -4.0458$$

$$f^{-1}(x) = \frac{10^{\left(\frac{x+5}{2}\right)} - 3}{2}$$

$$0 = \frac{10^{\left(\frac{x+5}{2}\right)} - 3}{2} \quad 0 = 10^{\left(\frac{x+5}{2}\right)} - 3 \quad 10^{\left(\frac{x+5}{2}\right)} = 3 \quad \frac{x+5}{2} = \log 3 \quad x = 2\log 3 - 5 \approx -4.0458$$

$$y = \frac{10^{\left(\frac{0+5}{2}\right)} - 3}{2} \quad y = \frac{10^{\left(\frac{5}{2}\right)} - 3}{2} \approx 156.61$$

Measuring Earthquakes

Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

On the Richter scale, the magnitude R of an earthquake of intensity I is given by $R = \frac{\ln I - \ln I_0}{\ln 10}$ where I_0 is the minimum intensity used for comparison.

Write this as a single common logarithmic expression.

Using the change of base formula.

$$R = \frac{\ln I - \ln I_0}{\ln 10} = \frac{\frac{\log I}{\log e} - \frac{\log I_0}{\log e}}{\frac{\log 10}{\log e}} = \frac{\log I - \log I_0}{\log 10} = \frac{\log I - \log I_0}{1} = \log \frac{I}{I_0}$$

Another Look

Use the product rule, the quotient rule, and the power rule, to expand and condense logarithmic expressions.

On the Richter scale, the magnitude R of an earthquake of intensity I is given by $R = \frac{\ln I - \ln I_0}{\ln 10}$ where I_0 is the minimum intensity used for comparison.

Write this as a single common logarithmic expression.

Using the converse of the change of base formula.

$$R = \frac{\ln I - \ln I_0}{\ln 10} = \frac{\ln I}{\ln 10} - \frac{\ln I_0}{\ln 10} = \log I - \log I_0 = \log \frac{I}{I_0}$$